

Noromectin

1.87% Oral Paste for Horses

Types of worms that affect horses


Pinworms

Diagnosis:

Tail base irritation with white eggs possibly visible around the peri-anal region

Treatment:

Adults- Benzimidazoles, Pyrantel or Macrocytic Lactones such as Ivermectin or Moxidectin



Small red worms


Diagnosis:

Faecal egg count to show presence of adult worms

Treatment:

Adults- Benzimidazoles, Pyrantel or Macrocytic Lactones such as Ivermectin or Moxidectin

(Encysted Small Redworm Larvae: Moxidectin or 5 day course of Benzimidazoles)



Bots

Diagnosis:

Bot larvae over-winter attached to the lining of the stomach

Treatment:

Macrocytic Lactones such as Ivermectin or Moxidectin




Ascarids

Diagnosis:

Faecal egg count

Treatment:

Adults and Larvae- Pyrantel or Macrocytic Lactone such as Ivermectin or Moxidectin




Large red worms

Diagnosis:

Faecal egg count

Treatment:

Adults and larvae- Benzimidazoles or Macrocytic Lactones such as Ivermectin or Moxidectin




Lung worms

Diagnosis:

Uncommon in horses

Treatment:

Macrocytic Lactones such as Ivermectin or Moxidectin



When to worm with Noromectin

1.87% Oral Paste for Horses

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Noromectin	BOTS	Routine roundworm treatment									BOTS	BOTS
Small encysted red worm												
Tapeworm												

What does Noromectin treat compared to other ingredients?

	Group one		Group two	Group three	Group four
	Macrocyclic lactones		Benzimidazoles	Pyrantel Ebonate	Praziquantel
	Ivermectin Noromectin	Moxidectin			
CESTODES-(tapeworm)					
A.perfoliata				X (DOUBLE DOSE)	X
A.magna					X
P. mamillana					X
NEMATODES-(roundworms)					
Large Round Worm					
Adults	X	X	X	X	
Larvae	X	X	X		
Small Round Worm					
Adults	X	X	X	X	
Non mucosal larvae	X	X	X		
Encysted cyathostomes		X	X (5 DAY COURSE)		
Benzimidazole resistant strains	X	X		X	
Lungworm	X	X			
Pinworm					
Adult	X	X	X	X	
Larvae	X	X	X		
Ascarids					
Adult	X	X	X	X	
Larvae	X	X	X		
Hairworms	X	X			
Intestinal threadworms	X	X			
Neck threadworms	X	X			
INSECTA					
Bots	X	X			

NOROMECTIN - The Facts

Dial a dose syringe for easy dosing and administration.

Small and easy to handle syringe.

One syringe can treat horses up to 700kgs

(Treats two ponies weighing less than 350kgs each.

Or variants of, dependent on weight).

Does not kill tapeworm (praziquantel) or small encysted red worm (moxidectin).

In foals Noromectin can be given from 6-8 weeks of age.

For pregnancy and lactation, Noromectin can be given at any stage.

Packaging quantities: Syringe containing 7.49g of product in cartons of 1, 2 and 10 syringes.

NO flavouring.

Legal Category UK: POM-VPS ROI: LM

Manufactured and distributed in NI by: Norbrook Laboratories Ltd, Station Works, Newry, Co. Down, BT35 6JP.

Distributed in GB by: Norbrook Laboratories (GB) Ltd, 1 Saxon Way East, Oakley Hay Industrial Estate, Corby, NN18 9EX.

Distributed in ROI by: Norbrook Laboratories (Ireland) Limited, Rossmore Industrial Estate, Monaghan, County Monaghan.

Noromectin

1.87% Oral Paste for Horses

