

**PRESCRIPTION ANIMAL REMEDY  
KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY**

## **BETAMOX Injection**

### **ACTIVE CONSTITUENT:**

BETAMOX Injection is an off-white sterile suspension containing 150 mg/mL amoxicillin as amoxicillin trihydrate.

### **INDICATIONS:**

BETAMOX Injection is a broad spectrum semi-synthetic penicillin, bactericidal in action. *In vitro* it is effective against a wide range of amoxicillin sensitive Gram-positive and Gram-negative pathogenic bacteria which include; *Actinobacillus equuli*, *Antinobacillus lignieresii*, *Actinomyces bovis*, *Bacillus anthracis*, *Bordetella bronchiseptica*, *Clostridium* spp., *Corynebacterium* spp., *Erysipelothrix rhusiopathiae*, *Fusiformis* spp., *Haemophilus* spp., *Moraxella* spp., *Pasteurella* spp., *Proteus mirabilis*, Staphylococci and Streptococci in horses, cattle, pigs, sheep, dogs and cats.

### **DIRECTIONS FOR USE:**

#### **Restraints:**

**DO NOT USE** in lactating or pregnant ewes where milk or milk products may be used for human consumption.

#### **Precautions:**

BETAMOX Injection is not suitable for intravenous injection. Occasional local reaction may occur with use of BETAMOX Injection.

Amoxicillin like other penicillins, should not be used, "off-label", orally or parenterally in rabbits, hamsters, gerbils and guinea pigs.

Animals with functionally mature rumens should only be treated parenterally.

#### **Dosage and administration:**

**SHAKE WELL BEFORE USE.**

Administer by subcutaneous or intramuscular injection, then massage the injection site. Recommended dosage rate is 7 mg/kg bodyweight once a day for up to five days.

**Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may result in the need to extend the approved withholding period.**

Use the contents of the vial within 4 weeks of broaching and discard any unused portion.

## **WITHHOLDING PERIODS:**

### **MEAT:**

**DO NOT USE** less than 28 days before slaughter for human consumption.

### **MILK:**

**Cattle:** Milk collected from cows within 48 hours (4 milkings) following treatment **MUST NOT BE USED** for human consumption or processing, or fed to bobby calves.

**Sheep:** **DO NOT USE** in lactating or pregnant ewes where milk or milk products may be used for human consumption.

### **TRADE ADVICE:**

**EXPORT SLAUGHTER INTERVAL (ESI):** This product does not have an ESI established. For advice on the ESI, contact the manufacturer on 1800 665 866 before using this product.

### **FIRST AID:**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

### **STORAGE:**

Store below 25°C (Air-Conditioning). Do not freeze. Protect from light.

### **DISPOSAL:**

Dispose of empty container by wrapping with paper and putting in garbage.

### **FURTHER INFORMATION:**

After absorption, high levels of amoxicillin are found in kidney, urine, liver and bile. Amoxicillin has rapid bactericidal action.

Manufactured by: Norbrook Laboratories Limited, UK

Distributed by:

Norbrook Laboratories Australia Pty Limited

ACN 080 972 596

Unit 7/15- 21 Butler Way,

Tullamarine, VIC 3043

Free call: 1800 665 866

APVMA Approval No. 37049/1109

