

WARNING

**KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY**

Moxiclear

*For Puppies and Small Dogs up to 4 kg /
For Dogs 4-10 kg / For Dogs 10-25 kg /
For Dogs Over 25 kg*

Active Constituents:

100 g/L IMIDACLOPRID, 25 g/L MOXIDECTIN

For use on dogs and puppies over 7 weeks of age. One application lasts 4 weeks.

For the treatment and prevention of

- fleas (*Ctenocephalides* spp.); Larval flea stages are killed in the surroundings of MOXICLEAR treated pets. MOXICLEAR can be used as part of a treatment strategy for flea allergy dermatitis (FAD)

For the control of

- roundworms (*Toxocara canis*, *Toxascaris leonina*) including larval (L4) and immature stages;
- hookworms (*Ancylostoma caninum*, *Uncinaria stenocephala*), including larval (L4) and immature stages of all species
- whipworm (*Trichuris vulpis*)

For the treatment and control of

- ear mites (*Otodectes cynotis*)
- sarcoptic mange (*Sarcoptes scabiei*)

An aid in the treatment and control of demodectic mange (*Demodex canis*)

For the control of lice (*Trichodectes canis*, *Linognathus setosus*)

Net Contents:

Moxiclear For Puppies and Small Dogs up to 4 kg

[1x0.4 mL] [2x0.4 mL] [3x0.4 mL] [4x0.4 mL] [6x0.4 mL] pipette(s)

Moxiclear For Dogs 4-10 kg

[1x1.0 mL] [2x1.0 mL] [3x1.0 mL] [4x1.0 mL] [6x1.0 mL] pipette(s)

Moxiclear For Dogs 10-25 kg

[1x2.5 mL] [2x2.5 mL] [3x2.5 mL] [4x2.5 mL] [6x2.5 mL] pipette(s)

Moxiclear For Dogs Over 25 kg

[1x4.0 mL] [2x4.0 mL] [3x4.0 mL] [4x4.0 mL] [6x4.0 mL] pipette(s)

DIRECTIONS FOR USE:

Consult a veterinarian before use.

For external application only.

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain or distress when using this product other than as directed on the label.

DOSAGE

For puppies and small dogs up to 4 kg bodyweight	Use 1 PIPETTE of MOXICLEAR FOR PUPPIES AND SMALL DOGS UP TO 4 KG
For dogs 4-10 kg bodyweight	Use 1 PIPETTE of MOXICLEAR FOR DOGS 4-10 KG
For dogs 10-25 kg bodyweight	Use 1 PIPETTE of MOXICLEAR FOR DOGS 10-25 KG
For dogs over 25 kg bodyweight	Use 1 PIPETTE of MOXICLEAR FOR DOGS OVER 25 KG

For very large dogs over 50 kg, apply 2 pipettes of MOXICLEAR for Dogs Over 25 kg by dividing the dose between the shoulder blades, on the midline of the back between the hips, and at one point in between.

1. Remove the pipette from the outer sachet using scissors or fold along diagonal line to expose nick; tear back at nick.
2. Hold the pipette upright.
3. Tap the narrow part of the pipette to ensure the contents remains within the main body of the pipette.
4. Twist or snap back the tip.
5. Site of Application.

Dogs under 25 kg	Dogs 25-50 kg
1 PIPETTE	1 PIPETTE
Apply between the shoulder blades	Apply by dividing the dose between the shoulder blades, on the midline of the back between the hips, and at one point in between.

6. The animal should be standing for easy application. Part the hair until the skin is visible. Place the tip of the pipette on the skin and squeeze the pipette firmly twice to apply the solution directly on the skin. Avoid pet's eyes or mouth.
7. Record treatment date and repeat monthly or as directed.
8. The solvent in MOXICLEAR may mark certain materials including plastics, leather, fabrics and finished surfaces. Allow application site to dry before permitting contact with such materials.

GENERAL INSTRUCTIONS

FLEA CONTROL AND PREVENTION

MOXICLEAR has been shown experimentally to kill adult and larval fleas within 20 minutes. Death can occur due to contact by absorption via the flea intersegmental membranes. Stops fleas from feeding in 3-5 minutes. Fleas are not required to bite to receive a lethal dose.

MOXICLEAR can be used as part of a treatment strategy for flea allergy dermatitis (FAD). MOXICLEAR breaks the FAD cycle by eliminating fleas which are the cause. Animals with severe FAD may require additional veterinary treatment.

It is recommended that all dogs in the same environment should be protected with MOXICLEAR for dogs. Additionally all cats should be protected with MOXICLEAR for cats. MOXICLEAR may be used as the only product for flea control as it breaks the life cycle and kills fleas emerging from the environment to reinfest the dog. New fleas may continue to emerge from the environment for six weeks or longer (depending on climate) after treatment is initiated.

More rapid resolution of heavy environmental infestations may be obtained by additional environmental control measures using a suitable registered product. Exposure of dogs to water or shampooing 4 hours after treatment does not reduce the effectiveness of MOXICLEAR. Frequent swimming or washing can shorten the duration of flea control.

WORM CONTROL

MOXICLEAR controls the major gastrointestinal nematodes of dogs but does not control tapeworms. In order to control tapeworms use specific agent. Dogs can become reinfested with worms but regular use of MOXICLEAR will remove them.

If worm problems persist consult a veterinarian.

TREATMENT AND CONTROL OF EAR MITES

For the elimination of ear mites (*Otodectes cynotis*), MOXICLEAR should be applied once and is highly efficacious as a single dose. Animals not responding to the initial treatment may respond to a second monthly treatment. Such non responding animals should be monitored and veterinary advice sought if treatment is not successful. Monthly application of MOXICLEAR will treat any subsequent ear mite infestation. To maximise efficacy gentle cleaning of the ear canal is recommended.

TREATMENT AND CONTROL OF SARCOPTIC MANGE

MOXICLEAR has been shown to have good efficacy against sarcoptic mites (*Sarcoptes scabiei*) after a single dose. A second dose of MOXICLEAR a month later will result in continued improvement of the clinical signs associated with sarcoptic mange. Monthly application of MOXICLEAR will control any subsequent sarcoptic mite infestation.

AN AID IN THE TREATMENT AND CONTROL OF GENERALISED DEMODICOSIS

MOXICLEAR may be effective in the treatment of generalised demodicosis when administered at monthly intervals for 2 to 4 treatments.

LICE CONTROL

A single treatment of MOXICLEAR will control lice on dogs.

PUPPIES

Puppies may be treated with MOXICLEAR For Puppies and Small Dogs up to 4 kg from 7 weeks of age. If puppies younger than 7 weeks require flea control they may be safely treated with an alternative imidacloprid only ectoparasiticide. When worming puppies prior to commencing MOXICLEAR, use a specific worming agent at 2, 3, 4, and 6 weeks of age. Once treatment with MOXICLEAR every month is commenced worms will be continuously controlled.

PRECAUTIONS

Do not apply MOXICLEAR FOR DOGS to cats.
Use with caution in sick, debilitated or underweight animals.

PREGNANCY AND LACTATION

No primary embryo-toxic, teratogenic or reproductive toxic effects have been observed with imidacloprid or moxidectin in laboratory species. Evidence suggests that no adverse effects are to be expected in pregnant and lactating dogs. However, the safety of MOXICLEAR has not been established during pregnancy and lactation.

ADVERSE EVENTS

In extremely rare cases, application site disorders such as hair loss, redness and itching may occur.

STORAGE

Store below 25°C and protect from light.

PRESENTATION

MOXICLEAR FOR	Contents	ACVM No:
PUPPIES AND SMALL DOGS UP TO 4KG	0.4 mL	A11874
DOGS 4-10 KG	1.0 mL	
DOGS 10-25 KG	2.5 mL	
DOGS OVER 25 KG	4.0 mL	

Registered pursuant to the ACVM Act 1997, No. A11874
See www.foodsafety.govt.nz for registration conditions.

Registered to and Distributed by:
Norbrook NZ Ltd
KPMG Centre, 18 Viaduct Harbour Avenue
AUCKLAND, Ph: 0800 224 022

READ LABEL BEFORE USE

If medical advice is needed, have product label at hand.

WARNING



Harmful if swallowed. May be harmful in contact with skin. Causes serious eye irritation. May cause an allergic skin reaction. Suspected of damaging fertility or the unborn child. May cause harm to breast-fed children. May cause damage to organs through prolonged or repeated exposure. Very toxic to aquatic life or Very toxic to aquatic life with long lasting effects. Toxic to the soil environment, terrestrial vertebrates and terrestrial invertebrates.

HANDLING PRECAUTIONS

Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Do not breathe vapours. Avoid contact during pregnancy/while nursing. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Contaminated work clothing should not be allowed out of the workplace. Avoid release to the environment. Wear protective gloves/eye protection. Use personal protective equipment as required. Collect spillage. Store locked up.

FIRST AID

IF SWALLOWED: Call the National Poisons Centre 0800 POISON (0800 764766) or a doctor/physician if you feel unwell. Rinse mouth.
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.
IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs: Get medical advice/attention. Wash contaminated clothing before reuse. **IF exposed or concerned:** Get medical advice/attention. Get medical advice/attention if you feel unwell.

DISPOSAL

Dispose of used pipettes by wrapping in paper and placing in garbage. Do not empty into waterways or stormwater.



Norbrook[®]