

WARNING
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

Moxiclear

*For Kittens and Small Cats up to 4 kg /
For Cats Over 4 kg*

Active Constituents:
100 g/L IMIDACLOPRID, 10 g/L MOXIDECTIN

For use on cats and kittens from 9 weeks of age. One application lasts 4 weeks.

For the treatment and prevention of

- fleas (*Ctenocephalides* spp.); Larval flea stages are killed in the surroundings of MOXICLEAR treated pets. MOXICLEAR can be used as part of a treatment strategy for flea allergy dermatitis (FAD).

For the control of

- roundworms (*Toxocara cati*, *Toxascaris leonina*) including larval (L4) and immature adult stages
- hookworms (*Uncinaria stenocephala**), including larval (L4) and immature stages of all species;

For the treatment and control of

- lungworm (*Aelurostrongylus abstrusus*)
- ear mites (*Otodectes cynotis*)

* not clinically significant in New Zealand

Net Contents:

Moxiclear For Kittens and Small Cats up to 4 kg
[1x0.4 mL] [2x0.4 mL][3x0.4 mL] [4x0.4 mL][6x0.4 mL] pipette(s)

Moxiclear For Cats Over 4 kg
[1x0.8 mL] [2x0.8 mL][3x0.8 mL] [4x0.8 mL] [6x0.8 mL] pipette(s)

DIRECTIONS FOR USE:

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain or distress when using this product other than as directed on the label. For external application only.

DOSAGE

For kittens and small cats up to 4 kg bodyweight	Use 1 PIPETTE of MOXICLEAR FOR KITTENS AND SMALL CATS UP TO 4 KG
For cats over 4 kg bodyweight	Use 1 PIPETTE of MOXICLEAR FOR CATS OVER 4 KG

1. Remove the pipette from the outer sachet using scissors or fold along diagonal line to expose nick; tear back at nick.
2. Hold the pipette upright.
3. Tap the narrow part of the pipette to ensure the contents remains within the main body of the pipette.
4. Twist or snap back the tip.
5. Site of Application.

Kittens and small cats up to 4 kg	Cats over 4 kg
1 PIPETTE	1 PIPETTE
Apply on the back of the head	Apply on the back of the head

6. The animal should be standing for easy application. Part the hair until the skin is visible. Place the tip of the pipette on the skin and squeeze the pipette firmly **twice to apply the solution directly on the skin**. Avoid pet's eyes or mouth. On very rare occasions skin reactions such as hair loss, redness and itching may occur. This is usually a single occurrence and hair will grow back uneventfully.
7. Record treatment date and repeat monthly or as directed.
8. The solvent in MOXICLEAR may mark certain materials including plastics, leather, fabrics and finished surfaces. Allow application site to dry before permitting contact with such materials.

GENERAL INSTRUCTIONS

FLEA CONTROL AND PREVENTION

MOXICLEAR has been shown experimentally to kill adult and larval fleas within 20 minutes. Death can occur due to contact by absorption via the flea intersegmental membranes. Stops fleas from feeding in 3-5 minutes. Fleas are not required to bite to receive a lethal dose.

MOXICLEAR can be used as part of a treatment strategy for flea allergy dermatitis (FAD). MOXICLEAR breaks the FAD cycle by eliminating fleas which are the cause. Animals with severe FAD may require additional veterinary treatment.

It is recommended that all cats in the same environment should be protected by MOXICLEAR for cats and all dogs with MOXICLEAR for dogs. MOXICLEAR may be used as the only product for flea control as it breaks the life cycle and kills fleas emerging from the environment to reinfest the cat. New fleas may continue to emerge from the environment for six weeks or longer (depending on climate) after treatment is initiated. More rapid resolution of heavy environmental infestations may be obtained by additional environmental control measures using a suitable registered product.

WORM CONTROL

MOXICLEAR controls the major gastrointestinal nematodes of cats but does not control tapeworms. In order to control these worms use specific agent. Cats can become reinfested with worms but regular use of MOXICLEAR will remove them.

If worm problems persist, consult a veterinarian.

TREATMENT AND CONTROL OF EAR MITES

For the elimination of ear mites (*Otodectes cynotis*), MOXICLEAR should be applied once and is highly efficacious as a single dose. Cats not responding to the initial treatment may respond to a second monthly treatment however this has not been established for MOXICLEAR. Such non responding animals should be monitored and veterinary advice sought if treatment is not successful. Monthly application of MOXICLEAR will treat any subsequent ear mite infestation. To maximise efficacy gentle cleaning of the ear canal is recommended.

TREATMENT AND CONTROL OF LUNGWORM

For the treatment and control of lungworm (*Aelurostrongylus abstrusus*), MOXICLEAR is highly efficacious as a single dose. Monthly application of MOXICLEAR will control any subsequent lungworm infections.

KITTENS

Kittens may be treated with MOXICLEAR FOR KITTENS AND SMALL CATS UP TO 4 KG from 9 weeks of age. If kittens younger than 9 weeks require flea control they may be safely treated with an alternative imidacloprid only ectoparasiticide.

For control of roundworms and hookworms kittens require worming at 4, 6 and 8 weeks of age. When worming kittens prior to commencing MOXICLEAR, use a specific worming agent. Once treatment with MOXICLEAR every month is commenced worms will be continuously controlled.

PRECAUTIONS

Use with caution in sick, debilitated or underweight animals. Take care to avoid the product coming into contact with the eyes or mouth of the animal. Ingestion by the recipient or in-contact animals should be prevented. Transient salivation, vomiting and pupil dilation may follow accidental oral ingestion.

PREGNANCY AND LACTATION

No primary embryo-toxic, teratogenic or reproductive toxic effects have been observed with imidacloprid or moxidectin in laboratory species. Evidence suggests that no adverse effects are to be expected in pregnant and lactating cats. However, the safety of MOXICLEAR has not been established during pregnancy and lactation.

ADVERSE EVENTS

In extremely rare cases, application site disorders such as hair loss, redness and itching may occur.

STORAGE

Store below 25°C and protect from light.

PRESENTATION

MOXICLEAR FOR	Contents	ACVM No:
KITTENS AND SMALL CATS UP TO 4 KG	0.4 mL	A11873
CATS OVER 4 KG	0.8 mL	

Registered pursuant to the ACVM Act 1997, No. A11873
See www.foodsafety.govt.nz for registration conditions.

Registered to and Distributed by:
Norbrook NZ Ltd
KPMG Centre, 18 Viaduct Harbour Avenue
AUCKLAND, Ph: 0800 224 022

READ LABEL BEFORE USE

If medical advice is needed, have product label at hand.

WARNING



Harmful if swallowed. May be harmful in contact with skin. Causes serious eye irritation. May cause an allergic skin reaction. Suspected of damaging fertility or the unborn child. May cause harm to breast-fed children. May cause damage to organs through prolonged or repeated exposure. Very toxic to aquatic life or Very toxic to aquatic life with long lasting effects. Toxic to the soil environment, terrestrial vertebrates and terrestrial invertebrates.

HANDLING PRECAUTIONS

Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Do not breathe vapours. Avoid contact during pregnancy/while nursing. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Contaminated work clothing should not be allowed out of the workplace. Avoid release to the environment. Wear protective gloves/eye protection. Use personal protective equipment as required. Collect spillage. Store locked up.

FIRST AID

IF SWALLOWED: Call the National Poisons Centre 0800 POISON (0800 764766) or doctor/physician if you feel unwell. Rinse mouth.
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.
IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs: Get medical advice/attention. Wash contaminated clothing before reuse. IF exposed or concerned: Get medical advice/attention. Get medical advice/attention if you feel unwell.

DISPOSAL

Dispose of used pipettes by wrapping in paper and placing in garbage. Do not empty into waterways or stormwater.



Norbrook[®]