

LIVER FLUKE RAPID ANTIBODY TEST for SHEEP










SCAN THE QR CODE TO WATCH THIS 2 MINUTE **HOW TO VIDEO BEFORE** FIRST USE.




For use by Vets, Farmers and Livestock Handlers




- Each test is a single use test. Dispose after use. Each lateral flow device can only be used once.
- Use within one hour of opening.
- Do not use after the expiry date.
- Store at room temperature (15° 25°C).
- The test is designed for use with blood, but can used with plasma or serum.

SPECIMEN COLLECTION AND HANDLING



Collect blood from inside ear (preferred sample site) or on top of the nose (where hair meets the nose).


If using ear, rub the inside of the ear for a few seconds to increase blood flow. If using the nose, dry the area first with a clean tissue.






Wait 15 secs for the blood droplet to appear (2) before taking sample.


Gently collect blood sample to fill the collection cup (3).






Remove the lateral flow device from the foil pouch and ensure that the desiccant sachet is orange. (If green, discard and use new device).








INTERPRETATION OF TEST RESULTS

After 10 minutes evaluate the lateral flow device for the formation of a control line (C). If there is no visible control line the test is invalid.

If a control line (C) is visible, use the reference card to match up the colour of the **test line (T)** and assign a score to the test line. If there is no visible (T) line after 10 minutes, the animal has tested negative. If there any visible (T) line results, discuss with your vet.



Test score indications: 0-1 = Negative 2 = Inconclusive 3-10 = Positive

